

Table 3: Antagonistic activity of *Trichoderma* species against pathogenic fungi.

Sr. No.	Treatments	% Antagonism					
		<i>At</i>	<i>Rs</i>	<i>An</i>	<i>Gc</i>	<i>Fo</i>	<i>Mp</i>
1	<i>T. viride</i>	73.3(86.7)	51.1(59.1)	44.4(50.2)	74.4(87.8)	55.5(64.2)	84.4(97.4)
2	<i>T. harzianum</i>	90.4(97.5)	38.8(42.6)	54.4(62.9)	61.1(71.5)	66.6(78.7)	83.3(96.4)
3	<i>T. konongii</i>	77.7(91.1)	67.7(79.5)	57.7(66.8)	66.7(78.6)	61.1(71.5)	55.5(64.2)
4	<i>T. pseudokoningii</i>	71.1(84.4)	44.4(50.2)	42.2(47.2)	72.2(85.5)	62.2(72.8)	81.1(94.7)
5	<i>T. virens</i>	74.4(87.8)	46.6(52.7)	44.4(50.2)	68.9(81.1)	70.0(82.7)	57.7(66.8)
6	Control (mm)	90					

At-*Alternaria alternata*, *Rs*- *Rhizoctonia solani*, *An*-*Aspergillus niger*, *Gc*-*Geotrichum candidum*, *Fo*-*Fusarium oxysporum* f. sp. *spinaciae*, *Mp*-*Macrophomina phaseolina*

