

Table 4 Evaluation of *Trichoderma* spp. against pathogenic fungi by dual culture using Bell's scale\*(R).

<i>Trichoderma</i> spp	Test Pathogens					
	<i>A.alternata</i>	<i>R.solani</i>	<i>A.niger</i>	<i>G.candidum</i>	<i>F.oxysporum</i>	<i>M.phaseolina</i>
<i>T. viride</i>	R <sub>2</sub>	R <sub>4</sub>	R <sub>4</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>2</sub>
<i>T. harzianum</i>	R <sub>1</sub>	R <sub>4</sub>	R <sub>3</sub>	R <sub>3</sub>	R <sub>3</sub>	R <sub>2</sub>
<i>T. koningii</i>	R <sub>2</sub>	R <sub>3</sub>	R <sub>3</sub>	R <sub>3</sub>	R <sub>3</sub>	R <sub>3</sub>
<i>T. pseudokoningii</i>	R <sub>2</sub>	R <sub>4</sub>	R <sub>4</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>2</sub>
<i>T. virens</i>	R <sub>2</sub>	R <sub>4</sub>	R <sub>4</sub>	R <sub>3</sub>	R <sub>2</sub>	R <sub>3</sub>

\*Degree of antagonism

R<sub>1</sub>=*Trichoderma* completely overgrew pathogens (100% over growth);

R<sub>2</sub>=*Trichoderma* overgrew at least two-third pathogens (75% over growth);

R<sub>3</sub>=*Trichoderma* colonizes on one half of the pathogens (50% over growth);

R<sub>4</sub>=*Trichoderma* and the pathogens contact point after inoculation;

R<sub>5</sub>= Pathogens overgrow bioagent – *Trichoderma*.

