

Table 3 Infestation data for the main hosts of *B. invadens* and *C. cosyra* in northern Ghana

Fruit species	Avg. no. fruits /sample	No. <i>B. invadens</i> /kg fruit*	No. <i>C. cosyra</i> /kg fruit*
Mango	4.0	60.1 ± 3.0 d	11.5 ± 1.7 a
Marula plum	25.5	51.3 ± 2.8 c	47.7 ± 3.0 d
Soursop	19.8	45.2 ± 3.1 b	24.2 ± 2.6 b
Tropical almond	30.6	65.5 ± 4.4 e	85.4 ± 10.0 f
Persimmon	59.4	65.0 ± 6.8 e	36.3 ± 2.9.1 c
Syncomore fig	35.0	42.5 ± 1.4 b	72.4 ± 7.2 e
Jujube	177.5	37.4 ± 2.0 a	122.0 ± 11.1 h
African peach	18.5	70.7 ± 9.6 f	143.0 ± 19.0 i
Shea nut	23.7	80.1 ± 11.0 g	90.6 ± 12.1 f
Green pepper	20.7	47.8 ± 2.4 bc	32.4 ± 2.7 c
Icacina	35.5	64.4 ± 4.4 e	111.7 ± 15.8 g
Albarillo	167.0	35.6 ± 2.0 a	128.8 ± 12.0 h

Note: \* Mean number ± standar error. Means with same letters within columns are not significantly different at p = 0.05